



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ing point," and thus the inter-focal distance of the real or imaginary foci, as the case may be, plainly cuts all the co-axial circles orthogonally.

Corollary 8. In the theorem, the hyperbola's asymptotes being also tangents, the circle through the points (say e, f, e', f') in which any two tangents cut them, is also one of the co-axial system having the minor axis for its radical axis. So that $Ce.Cf=CS^2=Ce'.Cf'$.

DEPARTMENTS.

SOLUTIONS OF PROBLEMS.

ALGEBRA.

NOTE ON SHORT METHODS IN ARITHMETICAL CALCULATIONS.

From time to time, we have sent to us for publication, "short cuts" and "lightning methods." Most of these are of no theoretical and little practical value, and hence cannot be given a place in the MONTHLY.

The short cuts below were sent to us by Mr. Charles H. Case of Chicago, who is now nearly 81 years old. He says the list, consisting of eighteen examples, was prepared for the students of Wheaton College, November, 1896. We publish a few of them because we have found some of them useful in practical computation, having used them for years.

Mr. Case says, "The examples given should be wrought without the use of more figures than are used in the same." The principles used may be found mainly in the algebraic formulae given below.

$$(a \div b)(b \div a) = 1; \quad (a+b)^2 = a^2 + 2ab + b^2; \quad (a+b)(a-b) = a^2 - b^2.$$

$$1. \quad (5\frac{1}{2})^2 = 30\frac{1}{4}; \quad (7\frac{1}{2})^2 = 56\frac{1}{4}; \quad (65)^2 = 4225; \quad (88)^2 = 7744; \quad 96^2 = 9216; \\ 36^2 = 1296; \quad 76^2 = 5776.$$

$$2. \quad 625^2 = \begin{array}{r} 360625 \\ 3 \\ \hline 390625 \end{array}; \quad 876^2 = 767376; \quad 2496^2 = 6230016.$$

$$3. \quad 99649964125^2 = 9928129699281296015625 \\ + 198562592 \\ + 249124910 \\ \hline 9930115350113787015625$$

$$4. \quad (2436)^2 = \begin{array}{r} 5761296 \\ 1728 \\ \hline 5934096 \end{array}; \quad 68 \times 132 = 8976; \quad 8852 \times 8948 = 79207696; \quad 868 \times$$

$$932 = 808976.$$

$$5. 48\frac{17}{29} \times 49\frac{12}{29} = 2400\frac{697}{841}; \quad 1295\frac{57}{93} \times 1296\frac{36}{93} = 1677615\frac{7343}{8649}.$$

$$6. 7464 \times 7536 = 56248704; \quad 88044 \times 87956 = 7744000000 - 1936.$$

$$7. 2777\frac{7}{9} \times 4166\frac{2}{3} \times 666\frac{2}{3} \times 54 \times 24 \times 52 \times 7692307\frac{9}{13} \times 625 \times 125 \times 56 \times 32 \times 1428571428571428\frac{4}{7} \times 2083\frac{1}{3} \times 48 \times 833\frac{1}{3} \times 3125 \times 68543764287590 = \dots''$$

These and eleven other even longer computations are carried out mentally by Mr. Case.

The principle, $a^2 = (a-b)(a+b) + b^2$, may be used in the squaring of any number, though it is not so readily used if the numbers consist of more than two digits. Thus,

$$87^2 = (87-3)(87+3) + 3^2 = 84 \times 90 + 9,$$

$$92^2 = (92-2)(92+2) + 2^2 = 90 \times 94 + 4.$$

This is the principle used in several of Mr. Case's calculations. Thus, $(5\frac{1}{2})^2 = (5\frac{1}{2} - \frac{1}{2})(5\frac{1}{2} + \frac{1}{2}) + (\frac{1}{2})^2 = 30\frac{1}{4}$. ED. F.

330. Proposed by R. D. CARMICHAEL, Princeton, N. J.

An important function in the Theory of Numbers is one defined thus: $f(x)=1$ when $x>0$, $f(x)=0$ when $x=0$, $f(x)=-1$ when $x<0$. Two analytic expressions for $f(x)$ are the following:

$$f(x) = \lim_{n \rightarrow \infty} x^{1/(2n-1)}, \quad n=1, 2, \dots; \quad f(x) = \lim_{n \rightarrow \infty} \frac{(x+1)^n - (x+1)^{-n}}{(x+1)^n + (x+1)^{-n}}, \quad x > -1.$$

It is required to find other non-trigonometric analytic expressions for this function. (There are several representations of $f(x)$ by means of trigonometric functions.)

Remark by the PROPOSER.

Professor F. H. Safford, of the University of Pennsylvania, has sent me the following expressions for the function defined in the problem:

$$\frac{2}{\pi} \int_0^\infty \frac{\sin xz}{z} dz, \quad \frac{2}{\pi} \int_0^\infty \frac{x dz}{x^2 + z^2}, \quad \text{Lim.}_{m \rightarrow +\infty} \frac{e^{xm} - e^{-xm}}{e^{xm} + e^{-xm}}.$$

333. Proposed by R. D. CARMICHAEL, Princeton University.

Sum the infinite series

$$\frac{1}{(m+1)^2} + \frac{(2m-1)}{(2m+1)^2} + \frac{(3m-1)^2}{(3m+1)^4} + \frac{(4m-1)^3}{(4m+1)^5} + \frac{(5m-1)^4}{(5m+1)^6} + \dots$$

[No solution of this problem has been received.]

334. Proposed by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

$$\text{Sum the series, } 2^n - n \cdot 2^{n-2} + \frac{n(n-3)}{2!} 2^{n-4} - \frac{n(n-4)(n-5)}{3!} 2^{n-6} \\ + \frac{n(n-5)(n-6)(n-7)}{4!} 2^{n-8} - \frac{n(n-6)(n-7)(n-8)(n-9)}{5!} 2^{n-10} + \dots$$